

DECOMPOSITION FACTORS OF D-MODULES ON HYPERPLANE CONFIGURATIONS IN GENERAL POSITION

TILAHUN ABEBAW AND RIKARD BØGVAD

ABSTRACT. Let $\alpha_1, \dots, \alpha_m$ be linear functions on $\mathbb{C}^n$ and $X = \mathbb{C}^n \setminus V(\alpha)$, where $\alpha = \prod_{i=1}^m \alpha_i$ and $V(\alpha) = \{p \in \mathbb{C}^n : \alpha(p) = 0\}$. The coordinate ring $\mathcal{O}_X = \mathbb{C}[x]_\alpha$ of X is a holonomic A_n -module, where A_n is the n -th Weyl algebra and since holonomic A_n -modules have finite length, $\mathcal{O}_X$ has finite length. We consider a "twisted" variant of this A_n -module which is also holonomic. Define M_α^β to be the free rank 1 $\mathbb{C}[x]_\alpha$ -module on the generator α^β (thought of as a multivalued function), where $\alpha^\beta = \alpha_1^{\beta_1} \dots \alpha_m^{\beta_m}$ and the multi-index $\beta = (\beta_1, \dots, \beta_m) \in \mathbb{C}^m$. It is straightforward to describe the decomposition factors of M_α^β , when the linear functions $\alpha_1, \dots, \alpha_m$ define a normal crossing hyperplane configuration, and we use this to give a sufficient criterion on β for the irreducibility of M_α^β , in terms of numerical data for a resolution of the singularities of $V(\alpha)$.

1. INTRODUCTION

Definition 1.1. Let $V \cong \mathbb{C}^n$ be a finite-dimensional complex vector space.

- (i) A (finite) hyperplane configuration $A = \{H_1, \dots, H_m\}$ in V is a finite set of affine hyperplanes $H_i = V(\alpha_i)$ in V , where $\alpha_i : V \rightarrow \mathbb{C}$ is a non-zero polynomial of degree one.
- (ii) A hyperplane configuration A in V is said to be in *general position*, if

$$\{H_{i_1}, \dots, H_{i_p}\} \subset A, p \leq n \implies \dim_{\mathbb{C}}(H_{i_1} \cap \dots \cap H_{i_p}) = n - p$$

and

$$\{H_{i_1}, \dots, H_{i_p}\} \subset A, p > n \implies H_{i_1} \cap \dots \cap H_{i_p} = \emptyset.$$

- (iii) A hyperplane configuration A is said to be a *normal crossing configuration*, if for any set of hyperplanes $H_{i_1}, \dots, H_{i_k}$ that contain p , there is an affine change of local coordinates at p to new coordinates $x_1, \dots, x_n$, such that $\alpha_{i_1} = x_1, \dots, \alpha_{i_k} = x_k$. In particular any point $p \in V$ is contained in at most $n = \dim V$ of the hyperplanes in A .

If A is a normal crossing configuration, then the divisor $\alpha = \prod_{i=1}^m \alpha_i$ is a normal crossings divisor (see [7]). Any configuration in general position is a normal crossing configuration. The intersections $H \neq \mathbb{C}^n$ of a subset of the hyperplanes in A are called the *flats* of the configuration.

Example 1.2. Let $V = \mathbb{C}^2$. Then a set of lines in $\mathbb{C}^2$ defines

- (i) a hyperplane configuration in general position if no two are parallel and no three meet at a point and ;
- (ii) a normal crossing configuration if no three meet at a point.

There is an extensive literature on hyperplane arrangements [13]. The problem studied in this note is the following. Let $A = \{H_i = V(\alpha_i), i = 1, \dots, m\}$, be a hyperplane configuration, and set $X = \mathbb{C}^n \setminus V(\alpha)$, where $\alpha = \prod_{i=1}^m \alpha_i$. The coordinate

ring $\mathcal{O}_X$ of X is the localization $\mathbb{C}[x]_\alpha := \mathbb{C}[x_1, \dots, x_n]_\alpha$ and this is a holonomic A_n -module, where A_n is the n -th Weyl algebra. Consider the following twisted variant of this A_n -module, that corresponds to the multivalued function $\alpha^\beta = \alpha_1^{\beta_1} \dots \alpha_m^{\beta_m}$.

Definition 1.3. The module M_α^β is (as a $\mathbb{C}[x]_\alpha$ -module) the free rank 1 $\mathbb{C}[x]_\alpha$ -module on the generator α^β , where $\beta = (\beta_1, \dots, \beta_m) \in \mathbb{C}^m$. It is furthermore an A_n -module, defining

$$\partial_j(\alpha^\beta) = \sum_{i=1}^m \beta_i \frac{\partial_j(\alpha_i)}{\alpha_i} \alpha^\beta$$

for $j = 1, 2, \dots, n$ and extending to an action of A_n on M_α^β .

The A_n -module M_α^β is holonomic and so has finite length. The decomposition factors have support on the flats associated to the configuration (Proposition 2.6). In continuation of [1], where we treated the case $n = 2$, and inspired by [5] where incidentally the case of $\beta \in \mathbb{Z}^m$ is treated, our main interest lies in finding the decomposition factors of M_α^β . If A is a normal crossing configuration, this is not difficult. In Theorem 3.2, we prove that in that case M_α^β has exactly one decomposition factor with support on a flat $H = \bigcap_{k=1}^r H_{i_k}$ iff the associated β_{i_k} , $k = 1, \dots, r$ are integers. Normal crossing configurations occur when the singularities of $V(\alpha)$ are resolved. Applied to a such a resolution, Theorem 3.2 together with the decomposition theorem for D-modules, then gives our main result Theorem 5.4: a sufficient criterion for M_α^β to be irreducible for an arbitrary hyperplane arrangement, in terms of numerical data of the resolution and β .

2. SUPPORT OF D-MODULES

2.1. Definition. Let X be a smooth algebraic variety. We denote by $\mathcal{D}_X$ the sheaf of differential operators on X . If $X = \mathbb{C}^n$ this is the same as the sheaf on $\mathbb{C}^n$ associated to the Weyl algebra A_n . If X is an affine variety, we will by abuse of notation, identify $\mathcal{O}_X$ with $\Gamma(X, \mathcal{O}_X)$ and an $\mathcal{O}_X$ -module sheaf with its global sections as a module over $\Gamma(X, \mathcal{O}_X)$. If X is an affine open subset of $\mathbb{C}^n$ defined by $0 \neq \alpha \in \mathbb{C}[x]$, then $\mathcal{O}_X = \mathbb{C}[x]_\alpha$ and $\mathcal{D}_X = \mathbb{C}[x]_\alpha \otimes_{\mathbb{C}[x]} A_n$.

A holonomic $\mathcal{D}_X$ -module M is in particular an $\mathcal{O}_X$ -module and as such may be shown to have support on a closed variety $Z = \text{Supp} M$ (see [2]). If $\mathcal{I} \subset \mathcal{O}_X$ is the ideal associated to Z , any local section of M is annihilated by a large enough power of $\mathcal{I}$, and Z is the minimal closed subset with this property.

We will see later that the support of the irreducible factors of M_α^β are intersections of hyperplanes.

Example 2.1. (i) Let $M_1 = \mathbb{C}[x, y]_{xy} / (\mathbb{C}[x, y]_x + \mathbb{C}[x, y]_y)$. Then

$$\text{Supp} M_1 = V(x, y) = (0, 0).$$

(ii) Let $M_2 = \mathbb{C}[x, y]_x / \mathbb{C}[x, y]$. Then $\text{Supp} M_2 = V(x) = \{(0, y) : y \in \mathbb{C}\}$.

(iii) Let $M_3 = \mathbb{C}[x, y]$. Then $\text{Supp} M_3 = V(0) = \mathbb{C}^2$.

2.2. Some Basic Properties. Let M be a $\mathcal{D}_X$ -module and $U \subset X$ be an open subset. Then define $M|_U =: \mathcal{O}_U \otimes_{\mathcal{O}_X} M$; this is a $\mathcal{D}_U$ -module.

Lemma 2.2. Let $U \subset X$ be an open subset.

- (i) $\text{Supp} M|_U = U \cap \text{Supp} M$.
- (ii) $M|_U = 0 \Leftrightarrow \text{Supp} M \subset X - U =: Z$.

Proof. (i) is clear by the above description. Let $\mathcal{I}$ be the ideal of a closed subvariety Z , and let $j: U \rightarrow X$ be the inclusion. For any $\mathcal{O}_X$ -module $\mathcal{M}$ there exists an exact sequence of $\mathcal{O}_X$ -modules

$$\Gamma_Z M \subset M \longrightarrow j_* M|_U, \quad (2.1)$$

where (locally) $\Gamma_Z M = \{m \in M : \exists r, \mathcal{I}^r m = 0\}$. If $M|_U = 0$, then $\Gamma_Z M = M$ and this proves (ii) in one direction. The other direction is immediate. $\square$

It is easy to see that localization preserves irreducibility of $\mathcal{D}_X$ -modules. (Cf. also Lemmas 2.1-2 of [9]; since the formulation there is different and without proof, we give one.)

Proposition 2.3. *If M is an irreducible $\mathcal{D}_X$ -module and $U \subset X$ an open subset, then $M|_U =: \mathcal{O}_U \otimes_{\mathcal{O}_X} M$ is an irreducible $\mathcal{D}_U$ -module.*

Proof. Let $j: U \rightarrow X$ be the inclusion and denote the adjunction in (2.1) by $\delta: M \rightarrow j_* M|_U$. Since M is irreducible δ is either 0 or an injection. In the first case M is Z -torsion and $M|_U = 0$, and the lemma is trivially true. So assume that δ is an injection, and that $K \subset M|_U$ is an irreducible sub $\mathcal{D}_U$ -module. Then we have an injection $\gamma: j_* K \rightarrow j_* M|_U$. The intersection $\delta(M) \cap \gamma(j_* K)$ is either 0 or $\delta(M)$. In the second case $\delta(M) \subset \gamma(j_* K)$. By restricting to U , and using that $\delta|_U$ is the identity and $\gamma|_U$ is the inclusion $K \subset M|_U$, we get that $M|_U = K$. Hence the lemma is true in this case too. The first case implies that $\gamma: j_* K \rightarrow j_* M|_U / \delta(M)$ is an inclusion, which implies that $j_* K$ is a Z -torsion module. Hence $K = (j_* K)|_U = 0$. $\square$

Since localization is an exact functor we have the following corollary. We will use the notations $DF(M)$ for the set of decomposition factors of M and $c(M)$ for the number of decomposition factors.

Corollary 2.4. (i) $DF(M|_U) = \{M_i \in DF(M) : \text{Supp } M_i \cap U \neq \emptyset\}$.
(ii) $c(M|_U) \leq c(M)$.

2.3. The support of decomposition factors of M_α^β . That the support of decomposition factors of the modules we study are intersections of hyperplanes is easily seen by general arguments, but it is also possible to give a direct argument, as we will now do. First we will do a reduction.

Let $V_1 = \cap_{i=1}^m H_i$, and assume that this affine space has positive dimension. Choose a vector space complement V_2 to the affine subspace V_1 , such that $\mathbb{C}^n \cong V_1 \oplus V_2$, and denote the restriction of a linear function α to V_2 by $\tilde{\alpha}$.

Lemma 2.5 (Reduction lemma). i) M_α^β is isomorphic to the exterior tensor product $M_{\tilde{\alpha}}^\beta \hat{\otimes} \mathcal{O}_{V_1}$, as A_n -modules.
ii) If N_i , $i = 1, \dots, r$ are the decomposition factors of $M_{\tilde{\alpha}}^\beta$, then $N_i \hat{\otimes} \mathcal{O}_{V_1}$, $i = 1, \dots, r$ are the decomposition factors of M_α^β .
iii) $\text{supp}(N_i \hat{\otimes} \mathcal{O}_{V_1}) = (\text{supp } N_i) \times V_1$, $i = 1, \dots, r$.

Proof. i) follows by an affine change of coordinates. By Corollary 2.5 of [1] a decomposition factor of M_α^β is the external tensor product of a decomposition factor of $M_{\tilde{\alpha}}^\beta$ with $\mathcal{O}_{V_1}$. This gives ii). Finally iii) is immediate. $\square$

Proposition 2.6. *Decomposition factors of M_α^β have support on intersections of hyperplanes.*

Proof. Use the notation $H_S = \{p \in \mathbb{C}^n : \alpha_i(p) = 0, i \in S\}$, where $S \subset \{1, 2, \dots, m\}$. By the lemma we can reduce to the case when V_1 is a point or $V_1 = \emptyset$. Make induction on the number m of hyperplanes; for $m = 0$ the statement is immediate from the irreducibility of $\mathbb{C}[x]$ as an A_n -module. Assume it is true for an arbitrary arrangement with m hyperplanes. Then we have to prove the following: if N is a decomposition factor of M_α^β , where $\alpha = \prod_{i=1}^{m+1} \alpha_i$, then $H := \text{Supp } N$ is an intersection of hyperplanes.

By our reduction, there exists a hyperplane H_j such $H \not\subseteq H_j$, except in the case where H equals $\cap_{i=1}^{m+1} H_i$, and so in particular is a hyperplane intersection, and there is nothing to prove. Let $U = \mathbb{C}^n \setminus H_j$. Then by Proposition 2.3, $N|_U$ is non-zero and so a decomposition factor of $M_\alpha^\beta|_U$. On U , α_j is invertible and by Lemma 2.9 below, $\gamma: M_\alpha^\beta|_U \cong M_{\bar{\alpha}}^{\bar{\beta}}|_U$, where $\bar{\alpha} = \prod_{i=1, i \neq j}^{m+1} \alpha_i$ and $\bar{\beta} = (\beta_1, \dots, \hat{\beta}_j, \dots, \beta_{m+1})$. There is, again by Proposition 2.3, a decomposition factor N_1 of $M_{\bar{\alpha}}^{\bar{\beta}}$, such that $N_1|_U = \gamma(N|_U)$. By induction, $\text{supp} N_1 = H_S$ is an intersection of hyperplanes. Since twisting modules by automorphisms as in Lemma 2.9 preserves support, $\text{supp} N \cap U = H_S \cap U$. Since N is irreducible, $\text{supp} N$ is an irreducible subvariety, as is H_S , and so $\text{supp} N = \overline{\text{supp} N \cap U} = \overline{H_S \cap U} = H_S$. This proves the induction step. $\square$

Definition 2.7. Suppose that θ is an automorphism of $\mathcal{D}_X$. If M is a $\mathcal{D}_X$ -module, θ^*M is defined to be the $\mathcal{D}_X$ -module which consists locally of the same sections as M , but on which (local sections of) $\mathcal{D}_X$ acts by θ : if $P \in \mathcal{D}_X$, $m \in \theta^*M$, then $Pm = \theta(P)m$.

The following Lemma is clear.

Lemma 2.8. Let $\theta: \mathcal{D}_X \rightarrow \mathcal{D}_X$ denote an automorphism that is the identity on $\mathcal{O}_X$. Suppose M has decomposition factors M_i , $i = 1, \dots, l$. Then θ^*M has decomposition factors θ^*M_i , $i = 1, \dots, l$. In particular $c(M) = c(\theta^*M)$. The support of θ^*M_i equals the support of M_i .

We apply this in the following situation.

Proposition 2.9. Suppose that $X = \mathbb{C}^n$, so that $\mathcal{O}_X = \mathbb{C}[x]$ and let $U = X - V(\alpha_1, \dots, \alpha_l)$. Then $c(M_\alpha^\beta|_U) = c(M_{\bar{\alpha}}^{\bar{\beta}}|_U)$, where $\bar{\alpha} = \alpha_{l+1} \dots \alpha_m$ and $\bar{\beta} = (\beta_{l+1}, \dots, \beta_m)$.

Proof. By the preceding lemma, it suffices to construct an automorphism $\theta: \mathcal{D}_U \rightarrow \mathcal{D}_U$ that is the identity on $\mathcal{O}_U$ and such that $\theta^*(M_\alpha^\beta|_U) \cong M_{\bar{\alpha}}^{\bar{\beta}}|_U$. Further it is by induction enough to assume that $U = X - V(\alpha_1)$. Put $\alpha^\beta = \alpha_1^{\beta_1} \bar{\alpha}^{\bar{\beta}}$, where $\bar{\alpha} = \alpha_2 \dots \alpha_m$ and $\bar{\beta} = (\beta_2, \dots, \beta_m)$. Then $M_\alpha^\beta|_U = \mathbb{C}[x]_\alpha \alpha_1^{\beta_1} \bar{\alpha}^{\bar{\beta}}$ where α_1 is invertible. Define $\theta: \mathcal{D}_U \rightarrow \mathcal{D}_U$ by:

$$\theta(D) = D + \frac{\beta_1 D(\alpha_1)}{\alpha_1}$$

for all $D \in \text{Der}_{\mathbb{C}}(\mathbb{C}[x]) \subset \mathcal{D}_U$ and $\theta(r) = r$ for all $r \in \mathcal{O}_U$, and extend this to an endomorphism of $\mathcal{D}_U$. This is the desired automorphism, since it is easily checked that it has an inverse, and that the map

$$\rho: \theta^*(M_{\bar{\alpha}}^{\bar{\beta}}|_U) \rightarrow M_\alpha^\beta|_U,$$

defined by $\rho(r\bar{\alpha}^{\bar{\beta}}) = r\alpha^\beta$, is a $\mathcal{D}_U$ -isomorphism. $\square$

3. DECOMPOSITION FACTORS ON NORMAL CROSSINGS

We will now describe the decomposition factors of M_α^β for a normal crossing hyperplane configuration.

Lemma 3.1. Let $A = \{H_1, \dots, H_m\}$ be a nonempty normal crossing hyperplane configuration in $\mathbb{C}^n$. If $\cap_{i=1}^m H_i \neq \emptyset$, then $m \leq n$ and the arrangement is central. Assume that $\cap_{i=1}^m H_i = \emptyset$. Then, for each intersection $H(\neq \mathbb{C}^n)$ of a subset of the hyperplanes in A ,

- (i) there exists at least one hyperplane H_i , such that $H \not\subseteq H_i$, and
- (ii) there is a unique subset $S = \{i_1, \dots, i_k\} \subset \{1, \dots, m\}$ such that $H = H_S = \cap_{k=1}^r H_{i_k}$. Furthermore $H_j \supset H \iff j \in S$.

The following theorem, together with the fact that all decomposition factors have support on intersections of hyperplanes (Lemma 2.6), gives the number of decomposition factors of M_α^β and their support.

Theorem 3.2. *Let $A = \{H_1, \dots, H_m\}$ be a normal crossing arrangement on $\mathbb{C}^n$. Let $H_S = H_{i_1} \cap \dots \cap H_{i_k}$ be a flat. Consider decomposition factors of M_α^β .*

- (i) *There is at most one decomposition factor with support on H_S .*
- (ii) *There is exactly one decomposition factor with support on H_S if and only if $\beta_{i_1}, \dots, \beta_{i_k} \in \mathbb{Z}$.*

Proof. By Lemma 2.5 we may assume that $\cap_{i=1}^m H_i$ is either a point or $\emptyset$. If this intersection is a point p then as stated in Lemma 3.1, the arrangement is central and $m \leq n$, and the theorem is easy to prove, see [1, Prop. 3.1]. (The idea is that by a basis change it may be assumed that $\alpha_i = x_i$, $i = 1, \dots, m$ are coordinates at p , and then M_α^β is isomorphic to the external tensor product of $M_{\alpha_i}^{\beta_i}$ $i = 1, \dots, m$ on $\mathbb{C}^1$.)

This gives the induction basis for an inductive proof of (i) and (ii) on the number of hyperplanes in the arrangement, and it also allows us to assume that the intersection of all the hyperplanes in our configurations is empty. Assume that (i) and (ii) are true for all normal crossing arrangements with m hyperplanes, and let $A = \{H_i\}_{i=1}^{m+1}$ be a normal crossing arrangement with $m+1$ hyperplanes.

- (i) For $H_S = \mathbb{C}^n$ the statement of (i) follows trivially, since M_α^β has rank 1 as a module over O_U , where $U = \mathbb{C}^n \setminus V(\alpha)$. Let $H_S \neq \mathbb{C}^n$ be a flat of A . By Lemma 3.1 there is a hyperplane, which we may assume is H_{m+1} , such that $H_S \not\subseteq H_{m+1}$. Let $U = \mathbb{C}^n \setminus H_{m+1}$. By Proposition 2.3 any decomposition factor F of M_α^β with $\text{Supp} F = H$, satisfies that the restriction $F|_U$ is a non-zero irreducible D_U -module. By the proof of Lemma 2.9 the D_U -module $M_\alpha^\beta|_U$ is isomorphic to the module $M_{\tilde{\alpha}}^{\tilde{\beta}}|_U$ twisted by an automorphism, where $\tilde{\alpha} = \alpha_1 \dots \alpha_m$ and $\tilde{\beta} = (\beta_1, \dots, \beta_m)$. The twisting does not change the number of decomposition factors nor their support. By the induction assumption and Proposition 2.3 $M_{\tilde{\alpha}}^{\tilde{\beta}}|_U$ has at most one decomposition factor on $H_S \cap U$. So the same is true of M_α^β and by induction (i) is proved.
- (ii) Similarly, we can use induction to prove one direction of (ii). Assume that (ii) is true for all normal crossing configurations with m hyperplanes, and that we have an arrangement $\{H_i\}_{i=1}^{m+1}$. Let $H = H_{j_1} \cap \dots \cap H_{j_s}$ be a flat such that $\beta_{j_1} \in \mathbb{C} \setminus \mathbb{Z}$. By Lemma 3.1 there exists i such that $H \not\subseteq H_i$. By (i) there exists at most one decomposition factor of M_α^β with support H . Assume that there exists a decomposition factor of M_α^β , say F , with $\text{Supp} F = H$. Let $U = \mathbb{C}^n \setminus H_i$. Then $F|_U$ is simple and $\text{Supp} F|_U = H \cap U$. Let $\tilde{\alpha} = \alpha_1 \dots \hat{\alpha}_i \dots \alpha_{m+1}$ and $\tilde{\beta} = (\beta_1, \dots, \hat{\beta}_i, \dots, \beta_{m+1})$. Again arguing as in the proof of (i), we get that $M_{\tilde{\alpha}}^{\tilde{\beta}}$ has a decomposition factor with support on H . But this is a contradiction to the induction assumption. Therefore there is no decomposition factor of M_α^β which has support on H .

The converse of (ii) may be proven directly. Without loss of generality assume that $H = \cap_{i=1}^k H_i$ such that $\beta_1, \dots, \beta_k \in \mathbb{Z}$. Since we have a normal crossings arrangement, we may, by affine base change, assume that $\alpha_i = x_i$, $i = 1, \dots, k$. Then $M_\alpha^\beta \cong \mathbb{C}[x]_\alpha \tilde{\alpha}^{\tilde{\beta}}$, where $\tilde{\alpha}^{\tilde{\beta}} = \alpha_{k+1}^{\beta_{k+1}} \dots \alpha_m^{\beta_m}$. We first prove that there is one subfactor of M_α^β with support on H . Let $N \subset \mathbb{C}[x]_\alpha$

be the vector space over $\mathbb{C}$ generated by the set

$$\left\{ \prod_{j=1}^n x_j^{r_j} \prod_{l=k+1}^m \alpha_l^{s_l} \tilde{\alpha}^{\tilde{\beta}} : r_j \geq 0 \text{ if } j = k+1, \dots, n \right\}.$$

As a $\mathbb{C}[x]$ -module this is isomorphic to the localization of $\mathbb{C}[x]_{x_1 \dots x_k}$ to the open subset $\mathbb{C}^n \setminus \cup_{j=k+1}^m H_j$. It is clearly an A_n -module. Further let $N^+ \subset N$ be the vector space over $\mathbb{C}$ generated by the set

$$\left\{ \prod_{i=1}^n x_j^{r_j} \prod_{l=k+1}^m \alpha_l^{s_l} \tilde{\alpha}^{\tilde{\beta}} : r_{k+1} \geq 0, \dots, r_n \geq 0 \text{ \& } \exists l: 1 \leq l \leq k \text{ \& } r_l \geq 0 \right\}$$

As a $\mathbb{C}[x]$ -module this is isomorphic to the localization of $\Sigma_{j=1}^k \mathbb{C}[x]_{x_1 \dots \hat{x}_j \dots x_k}$ to $\mathbb{C}^n \setminus \cup_{j=k+1}^m H_j$. Again it is clearly an A_n -module, and N/N^+ is nonzero, since the quotient

$$\mathbb{C}[x]_{x_1 \dots x_k} / (\Sigma_{j=1}^k \mathbb{C}[x]_{x_1 \dots \hat{x}_j \dots x_k})$$

is non-zero, with support on H , and $H \cap \mathbb{C}^n \setminus \cup_{j=k+1}^m H_j \neq \emptyset$, by Lemma 3.1 (ii). Then N/N^+ is a nonzero A_n -module, which is a decomposition factor of M_α^β . As noted above $\text{Supp}(N/N^+) = V(\alpha_1, \dots, \alpha_k) = \cap_{i=1}^k H_i = H$, and hence there must be at least one irreducible decomposition factor with support on H .

This completes the proof of the theorem. $\square$

It should be noted that the proof only uses that α_i such that $\alpha_i(p) = 0$ form part of a system of parameters at each point p , and hence works in this situation for an arbitrary smooth affine variety.

Definition 3.3. Let A be a hyperplane configuration in $\mathbb{C}^n$ and $\mathcal{H}$ be the set of all nonempty intersections of hyperplanes in A , including $\mathbb{C}^n$ itself considered as the intersection over the empty set. Define a relation $x \leq y$ in $\mathcal{H}$ if $x \supseteq y$ (as subsets of $\mathbb{C}^n$). In other words, $\mathcal{H}$ is partially ordered by reverse inclusion. We call $\mathcal{H}$ the intersection poset of A .

We can not give formulas for the cardinality $|\mathcal{H}|$ valid for all normal crossings arrangements, but in the special case when A is a hyperplane configuration in general position this is possible. In that case any intersection of less than or equal to n hyperplanes gives a flat (see [11]) :

$$|\mathcal{H}| = \sum_{k=0}^n \binom{m}{k}.$$

Corollary 3.4. Let $M_\alpha^\beta = \mathbb{C}[x]_\alpha \alpha^\beta$, where $\alpha^\beta = \alpha_1^{\beta_1} \dots \alpha_m^{\beta_m}$ and the set $\{\alpha_1, \dots, \alpha_m\}$ defines a hyperplane configuration in general position. Assume that $\beta_1, \dots, \beta_k \in \mathbb{Z}$ and $\beta_{k+1}, \dots, \beta_m \in \mathbb{C} \setminus \mathbb{Z}$. Then the number of decomposition factors of M_α^β , $c(M_\alpha^\beta)$ is given by:

$$c(M_\alpha^\beta) = \sum_{j=0}^n \binom{k}{j},$$

where $\binom{k}{j} = 0$ if $j > k$.

Proof. In Theorem 3.2 we proved that M_α^β has exactly one decomposition factor for each flat H , where $H = H_{i_1} \cap \dots \cap H_{i_s}$ and $\beta_{i_1}, \dots, \beta_{i_s} \in \mathbb{Z}$. The sum in the theorem clearly counts the number of those, keeping in mind that by the hypothesis on the arrangement, each non-empty subset $S \subset \{1, \dots, m\}$ corresponds to a unique flat. $\square$




FIGURE 1. A hyperplane configuration defined by the polynomial $\alpha = xy(x + y + 1)$ and the Hasse diagram of its intersection poset.

4. EXAMPLE

4.1. **The A_2 -module $M_\alpha^\beta = \mathbb{C}[x, y]_{xy(x+y+1)} x^{\beta_1} y^{\beta_2} (x + y + 1)^{\beta_3}$.** We will describe the decomposition factors of M_α^β for different cases of β_1, β_2 and β_3 , as an example for the previous section. The following Theorem summarizes the results.

Theorem 4.1. *Consider the A_2 -module $M_\alpha^\beta = \mathbb{C}[x, y]_{xy(x+y+1)} x^{\beta_1} y^{\beta_2} (x + y + 1)^{\beta_3}$.*

- (i) *If $\beta_1, \beta_2, \beta_3 \in \mathbb{C} \setminus \mathbb{Z}$, then $c(M_\alpha^\beta) = 1$ and hence M_α^β is irreducible.*
- (ii) *If $\beta_1 \in \mathbb{Z}$ and $\beta_2, \beta_3 \in \mathbb{C} \setminus \mathbb{Z}$, then $c(M_\alpha^\beta) = 2$.*
- (iii) *If $\beta_1, \beta_2 \in \mathbb{Z}$ and $\beta_3 \in \mathbb{C} \setminus \mathbb{Z}$, then $c(M_\alpha^\beta) = 4$.*
- (iv) *If $\beta_1, \beta_2, \beta_3 \in \mathbb{Z}$, then $c(M_\alpha^\beta) = 7$.*

Proof. The linear functions x, y and $x + y + 1$ define a general position hyperplane configuration in $\mathbb{C}^2$. Therefore, by the previous section, the number of the decomposition factors of M_α^β is the number of flats H such that the corresponding β' s of the linear functions of the hyperplanes defining H are integers.

- (i) If $\beta_1, \beta_2, \beta_3 \in \mathbb{C} \setminus \mathbb{Z}$, then $H = \mathbb{C}^2$, considered as the empty intersection, is the only flat such that the β' s of the linear functions defining the hyperplanes defining $\mathbb{C}^2$ are integers. Therefore M_α^β has only one decomposition factor with support on $\mathbb{C}^2$ and hence M_α^β is irreducible with support on $\mathbb{C}^2$.
- (ii) If $\beta_1 \in \mathbb{Z}$ and $\beta_2, \beta_3 \in \mathbb{C} \setminus \mathbb{Z}$, then $H_1 = \mathbb{C}^2$, considered as the empty intersection and $H_2 = V(x)$ are the flats with corresponding integer β' s. We can also see this by the following composition series:

$$\mathbb{C}[x, y]_{y(x+y+1)} y^{\beta_2} (x + y + 1)^{\beta_3} \subset \mathbb{C}[x, y]_{xy(x+y+1)} x^{\beta_1} y^{\beta_2} (x + y + 1)^{\beta_3}$$

where

$$\mathbb{C}[x, y]_{y(x+y+1)} y^{\beta_2} (x + y + 1)^{\beta_3}$$

and

$$\mathbb{C}[x, y]_{xy(x+y+1)} x^{\beta_1} y^{\beta_2} (x + y + 1)^{\beta_3} / \mathbb{C}[x, y]_{y(x+y+1)} y^{\beta_2} (x + y + 1)^{\beta_3}$$

are easily seen to be irreducible A_2 -modules. Therefore, as predicted, $c(M_\alpha^\beta) = 2$, with one decomposition factor with support on $V(x)$ and one decomposition factor with support on the whole space $\mathbb{C}^2$.

- (iii) If $\beta_1, \beta_2 \in \mathbb{Z}$ and $\beta_3 \in \mathbb{C} \setminus \mathbb{Z}$, then $H_1 = \mathbb{C}^2$, considered as the empty intersection and $H_2 = V(x), H_3 = V(y)$ and $H_4 = V(x, y)$ are the flats with

corresponding integer $\beta's$. Consider the following sequence of vector spaces over $\mathbb{C}$:

$$R_0 \subset R_1 \subset M_\alpha^\beta,$$

where $R_0 = \mathbb{C}[x, y]_{(x+y+1)}(x+y+1)^{\beta_3}$, and R_1 is generated by the set

$$\left\{ \frac{\alpha_j^{r_j}(x+y+1)^s}{\alpha_i^{s_i}(x+y+1)^{\beta_3}} : s_j \geq 0, s_i \geq 1, \text{ for } i, j = 1, 2 \right\}.$$

Therefore $c(M_\alpha^\beta) = 4$ with one decomposition factor with support on each H_2, H_3, H_4 , one decomposition factor with support on $\mathbb{C}^2$.

- (iv) If $\beta_1, \beta_2, \beta_3 \in \mathbb{Z}$, then $H_1 = \mathbb{C}^2$, considered as the empty intersection and $H_2 = V(x), H_3 = V(y), H_4 = V(x+y+1), H_5 = V(x, y), H_6 = V(x, x+y+1)$ and $H_7 = V(y, x+y+1)$ are the flats with their $\beta's$ of the linear functions defining the hyperplanes defining $H_1, \dots, H_7$ are integers. Therefore $c(M_\alpha^\beta) = 7$, with exactly one decomposition factor with support on each of the flats.

□

5. RESOLUTION OF SINGULARITIES

Let $X = \mathbb{C}^n \setminus V(\alpha)$, and $\pi: \tilde{\mathbb{C}}^n \rightarrow \mathbb{C}^n$ be a resolution of the singularities of $V(\alpha)$. Let $E = \bigcup_{i=1}^s E_i$, where E_i are irreducible, be the exceptional divisor, and $Z = \pi(E)$ the center, so that $\tilde{\mathbb{C}}^n \setminus E \cong \mathbb{C}^n \setminus Z$. We then have that

$$\text{Div} \pi^*(\alpha_i) = \tilde{H}_i + r_1^i E_1 + \dots + r_s^i E_s, \quad (5.1)$$

where $\tilde{H}_i$ is the proper transform of $H_i = V(\alpha_i)$ and $r_j^i \in \mathbb{Z}_+$. Hence in an open affine $U \subset \tilde{\mathbb{C}}^n$,

$$\pi^*(\alpha_i)|_U = (\tilde{\alpha}_i^U (\gamma_1^U)^{r_1^i} \dots (\gamma_s^U)^{r_s^i}) \in \mathcal{O}_U \quad (5.2)$$

where $\tilde{H}_i \cap U = V(\tilde{\alpha}_i^U)$ and $U \cap E_j = V(\gamma_j^U)$. Some of the intersections $\tilde{H}_i \cap U$ or $U \cap E_j$ may be empty, but we suppress this from notation, so as to not make it cumbersome. The properties of $\tilde{\mathbb{C}}^n$ that we need are

- i) that there is an affine cover $\{U_i \subset \tilde{\mathbb{C}}^n, i = 1, \dots, r\}$, where each $U = U_i$ is isomorphic to an open affine subvariety of finite type of $\mathbb{C}^n$
- ii) $\pi^{-1}V(\alpha)$ is a normal crossing divisor.

For these standard properties see e.g. [7], or the construction in the present case in [6]. We will use the preceding section to study the D-module pullback $\pi^* M_\alpha^\beta$. The last statement above implies that Theorem 3.2 applies to the restriction of this module to each U_i .

Lemma 5.1. *Let $U = U_i$, $i = 1, \dots, r$. Then $\pi^* M_\alpha^\beta|_U \cong M_\alpha^{\tilde{\beta}}$ as D_U -modules, where $\tilde{\alpha} = \tilde{\alpha}_1^U \dots \tilde{\alpha}_m^U \gamma_1^U \dots \gamma_s^U$. (In the product we only use the factors for which the corresponding intersections $\tilde{H}_i \cap U$ or $U \cap E_j$ are non-empty.) Further (with the same restriction)*

$$(\tilde{\beta}_1, \tilde{\beta}_2, \dots, \tilde{\beta}_{m+s}) = (\beta_1, \beta_2, \dots, \beta_m, \sum_{i=1}^m r_1^i \beta_i, \dots, \sum_{i=1}^m r_s^i \beta_i).$$

Using the last section, we immediately obtain the decomposition factors of $\pi^* M_\alpha^\beta$. Extend the notation by defining $\tilde{H}_{m+i} := E_i$, $i = 1, \dots, s$.

Corollary 5.2. *The module $\pi^*(M_\alpha^\beta)$ has exactly one decomposition factor for each flat $\tilde{H} = \tilde{H}_{i_1} \cap \dots \cap \tilde{H}_{i_k}$ such that $\tilde{\beta}_{i_1}, \dots, \tilde{\beta}_{i_k} \in \mathbb{Z}$.*

Proof. Since the U_i cover $\tilde{\mathbb{C}}^n$, there is to any decomposition factor N of $\pi^*(M_\alpha^\beta)$ a U_i such that $N|_{U_i} \neq 0$. This restriction has, by Proposition 2.6, support on an intersection $\tilde{H} \cap U_i = \tilde{H}_{i_1} \cap \dots \cap \tilde{H}_{i_k} \cap U_i$, and by Theorem 3.2 and the preceding Lemma, this implies that $\tilde{\beta}_{i_1}, \dots, \tilde{\beta}_{i_k} \in \mathbb{Z}$. Furthermore there is by Theorem 3.2 at most one. Conversely, assume that $\tilde{H} = \tilde{H}_{i_1} \cap \dots \cap \tilde{H}_{i_k}$ is such that $\tilde{\beta}_{i_1}, \dots, \tilde{\beta}_{i_k} \in \mathbb{Z}$. There is U_i such that $\tilde{H} \cap U_i \neq \emptyset$, and hence there is by Theorem 3.2 a decomposition factor of $\pi^*(M_\alpha^\beta)|_{U_i}$ with support on $\tilde{H} \cap U_i$. By Corollary 2.4 this means that there is a decomposition factor of $\pi^*(M_\alpha^\beta)$ with support on $\tilde{H}$. $\square$

Corollary 5.3. *The module $\pi^*(M_\alpha^\beta)$ is irreducible if and only if*

$$\beta_1, \dots, \beta_m, \sum_{i=1}^m r_i^1 \beta_i, \dots, \sum_{i=1}^m r_i^s \beta_i \in \mathbb{C} \setminus \mathbb{Z}.$$

We can use this and the decomposition theorem to get a sufficient criterion for when M_α^β is irreducible. We will use the decomposition theorem in the general form stated by Kashiwara [9] and recently proved by Mochizuki [10]. (Our module is regular, so our application actually could have referred to old versions of the decomposition theorem.)

Theorem 5.4. *Assume that r_j^i are defined by the property (5.1) of a resolution of the singularities of $V(\alpha)$. Then M_α^β is irreducible if $\beta_1, \dots, \beta_m, \sum_{i=1}^m r_i^1 \beta_i, \dots, \sum_{i=1}^m r_i^s \beta_i \in \mathbb{C} \setminus \mathbb{Z}$.*

Proof. Given the condition in the theorem, by Corollary 5.3, $\pi^*(M_\alpha^\beta)$ is irreducible. Consider now the following pullback diagram:

$$\begin{array}{ccc} \tilde{U} & \xrightarrow{j'} & \tilde{\mathbb{C}}^n \\ \pi|_{\tilde{U}} \downarrow & & \downarrow \pi \\ U & \xrightarrow{j} & \mathbb{C}^n \end{array} \quad (5.3)$$

where $U = \mathbb{C}^n \setminus V(\alpha)$ and j' and j are the respective inclusions. By definition $M_\alpha^\beta = j_*(M_{\alpha|U}^\beta)$. Since j is an affine map, this remains true interpreted in the derived category $D(D_{\mathbb{C}^n})$, where j_* now is the derived functor of $\mathcal{O}_{\mathbb{C}^n}$ -modules (which is also the direct D-module image j_+). Similarly j'_* and $\pi|_{\tilde{U}}^*$ are exact functors, the latter since $\pi|_{\tilde{U}}$ is an isomorphism. By Theorem 1.7.3 of [8], pullbacks behave nicely in the derived category $D(D_{\tilde{\mathbb{C}}^n})$, so

$$\pi^* M_\alpha^\beta = \pi^* j_*(M_{\alpha|U}^\beta) = j'_* \pi|_{\tilde{U}}^*(M_{\alpha|U}^\beta), \quad (5.4)$$

and consequently

$$H^r \pi^* M_\alpha^\beta = 0, \text{ if } r \neq 0. \quad (5.5)$$

Now, in the derived category,

$$\pi_+ \pi^*(M_\alpha^\beta) = \pi_+ j'_*(\pi|_{\tilde{U}}^*(M_{\alpha|U}^\beta)) = j_*(\pi|_{\tilde{U}})_+ \pi|_{\tilde{U}}^*(M_{\alpha|U}^\beta) = j_*(M_{\alpha|U}^\beta) = M_\alpha^\beta. \quad (5.6)$$

But by the Decomposition Theorem (see e.g. [10]), $H^0 \pi_+ \pi^*(M_\alpha^\beta)$ is semisimple. However M_α^β is indecomposable, since it is rank 1 over the generic point and torsion free as a $\mathbb{C}[x]$ -module, and hence it is irreducible. $\square$

6. PLANE CASE

In this section we will exemplify the preceding, by describing the pullback to a resolution of singularities of the A_2 -module $M_\alpha^\beta = \mathbb{C}[x, y]_\alpha \alpha^\beta$, where $\alpha = \prod_{i=1}^m \alpha_i$, $\alpha_1 = x$, $\alpha_2 = y$ and $\alpha_i = x + c_i y$ for all $i = 3, \dots, m$ and $c_i \neq c_j$ for all $i \neq j$ and $\beta = (\beta_1, \beta_2, \dots, \beta_m) \in \mathbb{C}^m$. We only need to blow up the origin. This is the locus:

$$\tilde{\mathbb{C}}^2 = \{(x, y), [W_0, W_1]\} : xW_1 = yW_0\} \subset \mathbb{C}^2 \times \mathbb{P}^1$$

together with the map

$$\pi : \tilde{\mathbb{C}}^2 \longrightarrow \mathbb{C}^2$$

which is the restriction of the projection of $\mathbb{C}^2 \times \mathbb{P}^1$ onto the first factor. Let $U_1 \subset \tilde{\mathbb{C}}^2$ be the open subset given by $W_0 \neq 0$. In terms of Euclidian coordinates, $w_1 = \frac{W_1}{W_0}$ and we can write:

$$U_1 = \{(x, y), (w_1)\} : xw_1 = y\} = \{(x, xw_1, w_1)\} \subset \mathbb{C}^2 \times \mathbb{C}^1.$$

From this description we see that $U_1 \cong \mathbb{C}^2$ with coordinates x, w_1 . The restriction $\pi|_{U_1}$, is given by:

$$\pi(x, w_1) = (x, xw_1).$$

Similarly let $U_2 \subset \tilde{\mathbb{C}}^2$ be given by $W_1 \neq 0$, and put $w_0 = \frac{W_0}{W_1}$. Then:

$$U_2 = \{(x, y), (w_0)\} : x = yw_0\} = \{(yw_0, y, w_0)\} \subset \mathbb{C}^2 \times \mathbb{C}^1.$$

So $U_2 \cong \mathbb{C}^2$ with coordinates y, w_0 , and the restriction $\pi|_{U_2}$, is given by:

$$\pi(y, w_0) = (yw_0, y).$$

Clearly $\tilde{\mathbb{C}}^2 = U_1 \cup U_2$ and hence $\{U_1, U_2\}$ is an affine cover of $\tilde{\mathbb{C}}^2$. The exceptional divisor E is given in the chart U_1 by $x = 0$, and in U_2 by $y = 0$. Furthermore in U_1

$$\pi^*(\alpha_1) = (x), \pi^*(\alpha_2) = (xw_1), \pi^*(\alpha_i) = (x(1 + c_i w_1)), \quad i = 3, \dots, m,$$

and in U_2

$$\pi^*(\alpha_1) = (yw_0), \pi^*(\alpha_2) = (y), \pi^*(\alpha_i) = (y(w_0 + c_i)), \quad i = 3, \dots, m.$$

Hence, $\text{Div} \pi^*(H_i) = \tilde{H}_i + E$, $i = 1, \dots, m$, (and $U_1 = \tilde{\mathbb{C}}^2 \setminus H_1$ and $U_2 = \tilde{\mathbb{C}}^2 \setminus H_2$.) This gives us the numerical description of the resolution that we need to apply Corollary 5.2.

Proposition 6.1. *Consider $\pi_2^*(M_\alpha^\beta)$. Let $I = \{i | \beta_i \in \mathbb{Z}\}$ contain k elements, and let $|\beta| = \sum_{i=1}^m \beta_i$.*

- (i) *If $|\beta| \in \mathbb{Z}$, then there is exactly one decomposition factor with support on each of $\mathbb{C}^2$, E , H_i , $i \in I$ and one on each intersection $H_i \cap E$, $i \in I$, and these are all decomposition factors. All together $2(k+1)$.*
- (ii) *If $|\beta| \in \mathbb{C} \setminus \mathbb{Z}$, then there is exactly one decomposition factor with support on $\mathbb{C}^2$, and one on each H_i , $i \in I$ only. All together $k+1$.*

We can compare the criterion for irreducibility of M_α^β that was obtained in Theorem 5.4, with what was proved in the plane case in [1] and see that in the plane it also gives a necessary criterion. We ignore whether this is true in general.

Corollary 6.2. *In the plane case, $\pi^*(M_\alpha^\beta)$ is irreducible if and only if M_α^β is irreducible if and only if $\beta_1, \dots, \beta_m, \sum_{i=1}^m \beta_i \in \mathbb{C} \setminus \mathbb{Z}$.*

Proof. See [1]. □




FIGURE 2. The hyperplane configuration before and after the blowup in one of the affine charts.

ACKNOWLEDGEMENTS

We gratefully acknowledge the encouragement we got while working on these topics from Demissu Gemed, Addis Ababa University and would like to dedicate this note to his memory. We also want to thank Jan-Erik Björk and Rolf Källström for interesting, useful and fun discussions. Finally the first author also acknowledges the economical support of ISP, Uppsala University.

REFERENCES

- [1] Abebaw, T. and Bøgvad, R. , *Decomposition of D-modules over a hyperplane arrangement in the plane*, arXiv:**math.AG/0812.1370** v1 (2008), to appear Arkiv f. Mat.
- [2] Björk, J.-E., *Rings of differential operators*, North-Holland Publishing Co., Amsterdam, 1979.
- [3] Björk, J.-E., *Analytic D-modules and applications*, Kluwer Academic Publishers Group, Dordrecht, 1993.
- [4] Coutinho, S. C., *A primer of algebraic D-modules*, Cambridge University Press, Cambridge, 1995.
- [5] De Concini, C. and Procesi, C., *The algebra of the Box-spline*, arXiv:**math.NA/0602019** v1 (2006) .
- [6] De Concini, C. and Procesi, C., *Topics in hyperplane arrangements, polytopes and box-splines*, book to be published.
- [7] Harris, J., *Algebraic geometry*, Springer-Verlag, New York, 1995.
- [8] Hotta, R., Takeuchi, K. and Tanisaki, T., *D-modules, perverse sheaves, and representation theory*, Progress in Mathematics, 236, Translated from the 1995 Japanese edition by Takeuchi, Birkhäuser Boston Inc., Boston, MA, 2008
- [9] Kashiwara, M., *Semisimple holonomic D-modules*, Topological field theory, primitive forms and related topics (Kyoto, 1996), Progr. Math., 160, pp. 267–271, Birkhäuser Boston, Boston, MA, 1998.
- [10] Mochizuki, T., *Wild harmonic bundles and wild pure twistor D-modules*, **arXiv:0803.1344v3**
- [11] Orlik, P. and Terao, H., *Arrangements of hyperplanes*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 300, Springer-Verlag, Berlin, 1992
- [12] Saito, M. and Sturmfels, B. and Takayama, N., *Gröbner deformations of hypergeometric differential equations*, Springer-Verlag, Berlin, 2000.
- [13] Stanley, R. P., *An introduction to hyperplane arrangements*, in Geometric combinatorics, IAS/Park City Math. Ser., 13, pp. 389–496, Amer. Math. Soc., Providence, RI, 2007,

DEPARTMENT OF MATHEMATICS, ADDIS ABABA UNIVERSITY AND STOCKHOLM UNIVERSITY
 E-mail address: **tabebaw@math.aau.edu.et**, **abebaw@math.su.se**,

DEPARTMENT OF MATHEMATICS, STOCKHOLM UNIVERSITY
E-mail address: rikard@math.su.se